



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

Schröeter was much beloved, and greatly appreciated for his high attainments by those who knew him best. To the scientific world his loss is well nigh irreparable.—J. C. A.

*Collinsia bicolor*.—While studying the genus *Collinsia* with reference to future revision, certain peculiarities of structure in the flower of *C. bicolor* have come to light which seem to be of sufficient significance to merit some mention in the GAZETTE.

At the point where the upper pair of stamens become free from the corolla, the wings of the filaments turn into the sac of the corolla. After continuing attached to the corolla by one edge for a little way they end in free tips which are somewhat bearded. These tips point out into the sac so that they almost meet over the rudimentary fifth stamen or gland at the base of the sac.

The conclusion arrived at after careful study of the question was that their function is to guard the nectar gland. The utility of such a device can readily be seen. The four stamens are declined toward the lower lip of the corolla and clasped by its middle lobe; so that the insect must enter by this path in order to reach the pollen. But the throat is so large and with such a wide opening that the insect could easily enter and reach the gland without coming in contact with the pollen, were it not for these guards, which effectually bar this road to the nectar.

The same phenomenon has been observed in milder form in *C. franciscana*. Both species are easily distinguished from all others by this peculiarity.—ALICE E. KEENER, *Herbarium Lake Forest University*.